

RESEARCH

Open Access



Systemic inhibition of 5-lipoxygenase by MK-886 exacerbates apical periodontitis bone loss in a mouse model

Igor Bassi Ferreira Petean¹, Alice Corrêa Silva-Sousa¹, Raquel Assed Bezerra da Silva², Marília Pacífico Lucisano¹, Léa Assed Bezerra da Silva¹, Guilherme Piedade Assed de Castro¹, Manoel Damião Sousa-Neto¹, Lúcia Helena Faccioli² and Francisco Wanderley Garcia Paula-Silva^{1,2*}

Abstract

Background To investigate if 5-LO selective inhibitor (MK-886) could be used for systemic treatment of experimentally induced apical periodontitis in a mouse model.

Methods Twenty-four C57BL/6 mice were used. After coronal opening, a solution containing *Escherichia coli* LPS (1.0 µg/µL) was inoculated into the root canals of the lower and upper right first molars (n = 72 teeth). After 30 days apical periodontitis was established, and the animals were treated with MK-886 (5 mg/kg), a 5-LO inhibitor, for 7 and 14 days. The tissues were removed for histopathological and histometric analyses, evaluation of osteoclast number and gene expression for receptor activator of nuclear factor kappa-B (*Tnfrsf11a*), receptor activator of nuclear factor kappa-B ligand (*Tnfrsf11b*), osteoprotegerin (*Tnfrsf11b*), tartrate-resistant acid phosphatase (*Acp5*), matrix metalloproteinase-9 (*Mmp9*), cathepsin K (*Ctsk*) and calcitonin receptor (*Calcr*). Statistical data analysis was performed using Kruskal Wallis followed by Dunn's tests ($\alpha = 0.05$).

Results Administration of MK-886 for 7 days exerted no effect on apical periodontitis progression compared to LPS inoculation without treatment ($p = 0.3549$), while treatment for 14 days exacerbated bone loss ($p < 0.0001$). Administration of MK-886 enhanced osteoclastogenesis signaling and osteoclast formation within 7 days ($p = 0.0005$), but exerted no effect at 14 days ($p > 0.9999$). After 7 days of treatment, MK-886 induced mRNA expression for *Acp5* ($p = 0.0001$), *Calcr* ($p = 0.0003$), *Mmp9* ($p = 0.0005$) and *Ctsk* ($p = 0.0008$), however no effect in those gene expression was observed after 14 days ($p > 0.05$).

Conclusion Systemic treatment with MK-886 exacerbated LPS-induced apical periodontitis in a mouse model.

Keywords Apical periodontitis, Bacterial lipopolysaccharide, 5-Lipoxygenase, MK-886, In vivo model

Background

Apical periodontitis (AP) is a localized immune response against endodontic infection, a sequel of the host defense response to the microbial challenge that cause dental and bone resorption of apical and periapical tissues [1–3]. Recently, a meta-analysis confirmed a high global prevalence of AP, with 52% of pooled samples worldwide reporting at least one tooth with AP [3]. Thus, reduction of the bacteria load and their toxins and allow the host

*Correspondence:

Francisco Wanderley Garcia Paula-Silva
franciscogarcia@forp.usp.br

¹ School of Dentistry of Ribeirão Preto, University of São Paulo, Avenida do Café, s/n, Ribeirão Preto, São Paulo, Brazil

² Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, São Paulo, Brazil



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

response to proceed with repair to complete healing of the periapical pathosis and the restoration of the function are the ultimate goals of non-surgical root canal therapy (NSRCT) [1, 2, 4]. However, fail of NSRCT may occur in up to 32% in cases of teeth with AP [5, 6].

Considering that AP is a multifactorial disease, to reduce failure in the NSRCT approaches that address microbiology, quality of treatment and host response particularities are needed [7]. Because AP is a host tissue inflammatory response to bacterial infection of the root canal and serves the purpose of restraining the progression of infection towards bone tissue, systemic therapy would be useful in cases of difficult healing apical periodontitis such as lesions in patients with comorbidities or dysbiosis. Thus, recent research investigates coadjuvant therapies to limit the AP development and establishment, or improve their healing rates after NSRCT, with promising results through the use of anti-inflammatory drugs and probiotics via systemic or topical administration, in order to control the inflammatory process during AP establishment and repair [8–11].

The inflammatory response in apical periodontitis involves the generation of eicosanoids, synthesized from the metabolization of arachidonic acid by cyclooxygenase (COX) or lipoxygenase (LO) enzymes [2, 12–15]. Leukotrienes (LT), a class of metabolites generated downstream of 5-LO and 5-LO activating protein (FLAP) [16] have been found in rat inflamed dental pulp and human AP [17–19]. The enzyme 5-lipoxygenase (5-LO) and its metabolites stimulate the formation and activity of osteoclasts in vitro [20, 21], however, 5-LO gene ablation resulted in increased bone loss in cases of apical periodontitis when root canals were infected with Gram-negative *Fusobacterium nucleatum* [22] or it had no impact on the enlargement of apical periodontitis, when root canals were infected with micro-organisms from oral cavity [15]. Bone metabolism is regulated by receptor activator of nuclear factor kappa-B (RANK), receptor activator of nuclear factor kappa-B ligand (RANKL) and osteoprotegerin (OPG). Resorption occurs when there is an imbalance in the expression of RANKL and OPG, due to inflammatory conditions that result in increase in RANKL activity and decrease in the regulatory activity of OPG [2, 23].

Although it is mainly used for asthma treatment, MK-886 is a 5-lipoxygenase activating protein inhibitor can be a pharmacological strategy for intervention in leukotriene biosynthesis to control inflammatory diseases [24, 25]. Because 5-LO inhibition with CJ-13610 attenuated inflammation and bone loss in lipopolysaccharide (LPS)-induced periodontal disease [26] and we previously found that short term inhibition of 5-LO resulted in reduced inflammatory mediator and RANKL synthesis

during AP development [12], we sought to investigate the effects of systemic administration of MK-886 for 7 and 14 days in mature AP. Since impairment of catabolic signaling could be a therapeutic measure to prevent bone loss in cases of apical periodontitis, the aim of this study was to investigate whether 5-LO inhibitor (MK-886) could be used as systemic treatment of experimentally induced apical periodontitis in mice. The null hypothesis of this study was that 5-LO inhibition would not prevent bone resorption in apical periodontitis.

Methods

Animals

This research was conducted in accordance with Brazilian Guideline for the Care and Use of Animals in Teaching or Scientific Research Activities regulated by the National Council for the Control of Animal experimentation (Law 11.794/2008), conformed to Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines and the animal study protocol was approved by the Ethical Committee at the University of São Paulo (IRB approval number 12.1.60.53.8). C57BL/6 male mice (n = 24) (*Mus musculus*), 8-week-old, weighting 18–20 g, were used for this study. The sample size was determined according to a previous study [10]. For the operative procedures, animals were anaesthetized i.m. with 10% ketamine hydrochloride (150 mg/kg; National Pharmaceutical Chemistry Union Agener S/A, Embu-Guaçu, SP, Brazil) and xylazine (7.5 mg/kg; Dopaser, Labs Calier S/A, Barcelona, Spain). Anaesthesia was sustained during the operative procedures throughout the experimental period, and the animals were monitored by a veterinarian throughout the entire experimental period.

LPS inoculation for induction of apical periodontitis

A surgical table with a device for mandibular retraction was used to keep the animals immobilized with mouth open for direct access to the molars. The mandibular and maxillary right first molars of each animal were used for LPS inoculation into the root canals. The first molars of the left side were not inoculated and served as the negative control group. Access cavity preparations were performed with 1011 spherical diamond burs (KG Sorensen Ind. Com. Ltda., Barueri, SP, Brazil) and 10 µL of LPS suspension (1.0 µg/µL) of *E. coli* 0127: B8 (L3129; Sigma-Aldrich Corp., St. Louis, MO, USA) were inoculated into the root canals of each tooth with use of an automatic micropipette [10–15]. Conventional glass ionomer cement (S.S. White Dental Articles Ltda, Rio de Janeiro, RJ, Brazil), manipulated according to manufacturer's instructions, was used for restoring the teeth. After LPS

inoculation, the animals were kept under supervision for thirty days, in order to establish the apical periodontitis.

Systemic drug treatment

After apical periodontitis was established, the animals were treated with MK-886 (Merck Millipore, Darmstadt, Germany). MK-886 was dissolved in ethanol (100 µL), according to manufacturer’s recommendations, diluted in distilled water (400 µL), and provided by gavage (0.5 mL, 5 mg/kg body weight) daily for time intervals of 7 and 14 days. Healthy teeth from animals on the opposite side that did not receive medication were used as negative control and teeth from group which the pulp chamber filled with LPS without MK-886 treatment was used as a positive control. The groups were determined by inoculation with LPS (n = 6 teeth for each experimental time interval), LPS + MK-886 (n = 6 teeth for each experimental time interval). At time intervals of 7 and 14 days after starting with treatment, the animals were euthanized by means of anesthetic overdose and the tissues were collected. The Fig. 1 is a visual representation of the group distribution and study design.

Histological preparation

The upper and lower jaws were dissected and removed using surgical scissors. The blocks containing tooth and bone were fixed in 10% buffered formalin for 24 h at room temperature and demineralized in 10% EDTA (Merck S.A. Chemical Industries, Rio de Janeiro, RJ, Brazil) for approximately 21 days. After demineralization, the blocks of specimens were submitted to routine histological processing, washed in running water for 24 h, dehydrated in increasing concentrations of alcohol, diaphanized in xylol, and embedded in paraffin. The blocks were sectioned longitudinally, in the bucco-lingual direction, to obtain slices with a thickness of 5 µm. Sections were stained with haematoxylin and eosin (HE) for histopathological and histometric evaluation and submitted to tartrate-resistant acid phosphatase (TRAP) enzyme assay, for enzyme histochemical analysis.

Morphometric analysis of AP size under light microscopy

Longitudinal sections of the teeth were obtained and stained by haematoxylin–eosin. The analysis was qualitatively performed by using conventional light microscopy (Zeiss Axio Imager, Carl Zeiss AG Light Microscopy, Göttingen, Germany). All the analyses were performed using videomicroscopy with the Zeiss AxioVision Software (Carl Zeiss AG Light Microscopy), in conjunction with the AxioCam MRc5 (Carl Zeiss AG Light Microscopy) microscope and camcorder, at 20× magnification, operated in fluorescence mode (excitation at 460–500 nm and emission at 512–542 nm). For each




Fig. 1 Flow diagram of the study design

specimen, the area of apical periodontitis was delineated and measured in mm² on sections representing the largest diameter of the lesion, using the ImageJ 1.50i software (NIH, Bethesda, MD). The delineation of the lesion excluded intact structures (periodontal ligament, cementum and alveolar bone), and included areas of resorption and inflammatory infiltrate, according to previously described parameters [10].”

Determination of the activity and presence of osteoclasts by enzyme histochemical analyses for TRAP

The analysis of activity and the number of positive osteoclasts were analyzed and determined by counting the number of multinucleate TRAP-positive cells in the resorption lacunae in direct contact with the alveolar bone around the area of apical periodontitis. The tissue sections were deparaffinized and incubated in a solution containing 8 mg of naphthol AS-MX di-sodium phosphate (Sigma-Aldrich) in 500 µL of NN-dimethyl-formamide. This was followed by the addition of 50 mL of a 0.2 mol buffer solution/L sodium acetate (pH 5.0) containing 70 mg of Fast Red ITR (Sigma-Aldrich). Next,

the sodium tartrate substrate dihydrate (50 mmol/L) was added to the solution and incubated at 37 °C for 12 h. After this, the sections were washed in distilled water and stained with hematoxylin. Other sections were incubated with a medium without substrate as a control.

Quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR)

The maxillary right first molars of three animals were used for the Experimental Group and their maxillary left first molars, for the Control Group. RNA extraction was performed with RNeasy Mini kit (RNeasy® Mini, Qiagen Inc., Valencia, CA, USA) and samples were treated with DNase I (RNase-Free DNase Set; Qiagen Inc.), according to manufacturer's protocol. RNA integrity was analyzed using 1% agarose electrophoresis and quantity was estimated in NanoDrop 1000 (Thermo Fisher Scientific Inc., Wilmington, DE, USA) at 230, 260 and 280 nm wavelengths.

From 2000 ng of total RNA using random primers (High Quality cDNA Reverse Transcriptase Kits, Applied Biosystems, Foster City, CA, USA) complementary DNA (cDNA) was synthesized. Aliquots of 2 µL of the total cDNA were amplified by qRT-PCR using primers for *Tnfrsf11a* (RANK; Mm00437135), *Tnfsf11* (RANKL; Mm00441906), *Tnfrsf11b* (OPG; Mm01205928), *Acp5* (Mm00475698), *Mmp9* (Mm00442991), *Ctsk* (Mm00484039) and *Calcr* (Mm01197736) (TaqMan® Gene Expression Assay, Applied Biosystems) in a StepOne Plus equipment (Applied Biosystems). *Gapdh* (Mm99999915) was used as reference gene. Duplicates were used for qRT-PCR reactions and amplification was done with denaturation at 95 °C for 2 min; followed by 40 cycles of 95 °C for 1 s and 60 °C for 20 s. Relative quantification was performed using the $\Delta\Delta C_t$ Method.

Statistical analysis

Data were analyzed using GraphPad Prism 8.0 (GraphPad, San Diego, CA) and presented non normal distribution, therefore statistical analyses were performed using nonparametric Kruskal Wallis followed by Dunn's tests ($\alpha=0.05$).

Results

Administration of MK-886 favored apical periodontitis bone loss

Thirty days after inoculation of LPS into the root canals, apical periodontitis was formed and was characterized by a wide periodontal ligament space, increased recruitment of inflammatory cells and bone loss. Treatment with MK-886 for 7 days had no effect on the progression of apical periodontitis when compared with LPS inoculation without any treatment ($p=0.3549$), while treatment

for 14 days significantly exacerbated bone loss in apical periodontitis ($p<0.0001$) (Fig. 2).

MK-886 treatment increased periapical osteoclastogenesis signaling and osteoclast formation

Significant increase in the number of osteoclasts was observed after LPS inoculation into root canals, when compared with healthy teeth ($p<0.0001$). Treatment with MK-886 for 7 days augmented osteoclast formation significantly ($p=0.0005$) but had no effect at 14 days when compared with LPS inoculation without any treatment ($p>0.9999$) (Fig. 3).

The effect of MK-886, with regard to the expression of RANK, RANKL and OPG revealed that this administration induced significantly higher RANK expression when compared with LPS inoculation without any treatment, after both at 7 and 14 days ($p=0.0027$ and $p=0.0014$, respectively). Furthermore, treatment for 7 days induced significant RANKL expression ($p=0.0278$) without effect at 14 days ($p=0.9877$); while MK-886 significantly upregulated OPG after both at 7 and 14 days ($p=0.0481$ and $p=0.0009$, respectively). Higher expression of RANKL over OPG (Ratio >1) was found at 7 days for LPS and LPS+MK-886 treatment ($p=0.0099$ and $p=0.0038$, respectively), without significant difference between them ($p=0.8462$). No catabolic over anabolic signaling was detected at 14 days (Fig. 4).

Treatment with MK-886 stimulated early expression of genes involved in bone resorption

After 7 days of treatment, MK-886 induced significant mRNA expression of the enzymes TRAP ($p=0.0001$), MMP-9 ($p=0.0005$), cathepsin K ($p=0.0008$) and for calcitonin receptor ($p=0.0003$). After 14 days, TRAP, MMP-9 and calcitonin receptor expression induced after inoculation of LPS into the root canals was not modulated by treatment with MK-886 ($p=0.9849$, $p=0.0746$ and $p=0.9185$, respectively). Cathepsin K mRNA expression did not change as a result of LPS inoculation alone and after treatment with MK-886 ($p=0.2095$ and $p=0.7240$, respectively) (Fig. 5).

Discussion

In this study it was found that treatment of apical periodontitis using 5-LO inhibitor MK-866 for 7 days enhanced osteoclast signaling and formation and treatment for 14 days increased bone loss and resulted in larger apical periodontitis lesions. The null hypothesis of this study was therefore rejected.

Previous investigations have reported that at 28 days, periapical bone resorption occurs after inoculation of 10 µL of LPS suspension into the root canals [10, 11]. Therefore, in the present study, the period of 30 days




Fig. 2 Apical periodontitis establishment and status after treatment with MK-886. **A** Photomicrographs of healthy teeth, obtained after LPS inoculation into root canals, or at time intervals of 7 and 14 days after treatment with and without MK-886. HE, original 20× magnification (scale bar = 100 µm). **B** Graphic representation of the measurement of area of apical periodontitis, in µm² in the different groups at 7 and 14 days (median and interquartile range). Dashed line indicates the area of a healthy periodontal ligament. * $p < 0.05$ compared with healthy periodontal ligament; # $p < 0.05$ compared with inoculation of LPS alone into the root canals; & $p < 0.05$ comparison between 7 and 14 days of the same treatment

after inoculation of LPS was investigated, which meant that apical periodontitis was established prior to the treatment with MK-886. Furthermore, data of the size and morphological aspects of the area of apical periodontitis were evaluated at 30 days, when tissue necrosis, thickening of the periodontal space and recruitment of inflammatory mediators were observed, confirming the establishment of apical periodontitis.

Regarding to the results, it has previously been demonstrated that 5-lipoxygenase inhibition plays a dual-role in osteoclastogenesis during bacteria-induced AP formation. Early on (7 and 14 days), osteoclastogenesis signaling was down-regulated by the inhibition of 5-LO, but in the long-term (21 and 28 days) inhibition failed to prevent synthesis of catabolic mediators that resulted in increased bone loss [12]. Here, treatment with 5-LO inhibitor in an already established LPS-induced AP lesion, exacerbated the catabolic effects on

bone. Although the experimental strategy was different from this study in which we used LPS inoculation to promote a mature and established apical periodontitis, the observed behavior in osteoclast signaling was similar to that observed previously [12]. MK-886 treatment stimulated RANKL production at 21 days which was stabilized at 28 days, after the establishment of a mature apical periodontitis developed after root canal contamination from oral environment. It is worth mentioning that the osteoclastogenesis signaling was not coincident with the presence of active osteoclasts, once this is a dynamic process. Thus, we observed that initially (7 days) there was an overexpression of these genes, that resulted in a more robust osteoclastogenesis and consequently, bone resorption, which tends to reach a limit, that reflects in the levels of expression of genes that regulates this process after 14 days of treatment.




Fig. 3 Osteoclast formation after treatment with MK-886. **A** Photomicrographs representative of enzyme histochemical findings for the tartrate-resistant acid phosphatase enzyme (TRAP) obtained from healthy teeth, after LPS inoculation into the root canals, or after 7 and 14 days of treatment with MK-886. Original magnification 20 × (scale bar = 100 µm). **B** Graphic representation of the number of tartrate-resistant acid phosphatase enzyme (TRAP) positive osteoclasts per lesion in the different groups at 7 and 14 days (median and interquartile range). Dashed line indicates number of TRAP + osteoclasts in healthy apical periodontal ligament. * $p < 0.05$ compared with healthy periodontal ligament; # $p < 0.05$ compared with inoculation of LPS alone into the root canals; & $p < 0.05$ comparison between 7 and 14 days of the same treatment

One possible explanation of these catabolic effects on bone after treatment with MK-886, relies on the crucial role of leukotrienes as efficient chemotactic agents, which promote polymorphonuclear aggregation and degranulation and stimulate leukocyte adherence to the endothelial wall for transmigration of inflammatory cells [27]. Inhibition of synthesis of 5-LO metabolites might have impaired the immune and inflammatory cell recruitment and response that resulted in augmented bone resorption. Indeed, an impaired host innate immune system and severe AP were found in an experimental 5-LO knockout

murine model [22]. Moreover, the 5-LO pathway plays a critical role in the stimulation of inflammatory mediator synthesis in polymicrobial-induced apical periodontitis in mice [15].

In vivo inhibition or ablation of 5-LO has been shown to have controversial effects on several bone diseases, i.e., this may have anabolic [12, 14, 22] or catabolic [25, 28] effects. In vitro, the mechanism involved in osteoclast formation induced by 5-LO has been demonstrated [28, 29]. Leukotrienes B₄ (LTB₄) and C₄ (LTC₄), metabolites of 5-LO, induce RANKL and osteoclast differentiation




Fig. 4 Detection of RANK/RANKL/OPG genes after treatment with MK-886. Relative expression of *Tnfrsf11a*, *Tnfrsf11* and *Tnfrsf11b* genes that encode RANK, RANKL and OPG, after inoculation of LPS into root canals and at 7 and 14 days of treatment with MK-886 (median and interquartile range). * $p < 0.05$ compared with healthy periodontal ligament; # $p < 0.05$ compared with inoculation of LPS into root canals; & $p < 0.05$ comparison between 7 and 14 days of the same treatment




Fig. 5 Detection of genes related to bone metabolism after treatment with MK-886. Relative expression of *Acp5*, *Mmp9*, *Ctsk* and *Calcr* genes that encode the tartrate-resistant acid phosphatase enzyme (TRAP), matrix-9 metalloproteinase, cathepsin K and the calcitonin receptor, after LPS inoculation into root canals, and at 7 and 14 days after treatment with MK-886 (median and interquartile range). * $p < 0.05$ compared with healthy periodontal ligament; # $p < 0.05$ compared with inoculation of LPS into the root canals; & $p < 0.05$ comparison between 7 and 14 days of the same treatment

of bone marrow-derived and RAW-264.7 macrophages [20, 21]. In a previous in vitro study, we found that 5-LO pathway is involved in the osteoclastic differentiation hematopoietic lineage cells and that exogenous addition of LTB₄, loaded in microspheres (MS) inhibited osteoclastogenesis induced by M-CSF and RANKL [30]. The mechanism involved induction of *MMP9* gene expression and inhibition of *CALCR* and *CTSK*, without changing *ACP5* [30]. Furthermore, 5-LO activity may be modulated by nitric oxide (NO) since prolonged exposure to lipopolysaccharide inhibits macrophage 5-LO metabolism by induction of NO synthesis [31]. In addition, increased generation of hydrogen peroxide (H₂O₂) and an inhibited production of NO were observed after in-vitro incubation of MK-886 with peritoneal macrophages, which have been related to the replication of infectious agents [32]. Therefore, one possible explanation for the divergent effects observed here could be the fact that LTB₄ enhances phagocytosis and killing of microorganisms, and its long-term inhibition might have impaired infection control resulting in increased bone loss.

With regards to undesirable systemic effects of 5-LO inhibitor administration, according to the literature these drugs are generally well-tolerated, and the majority of adverse events described were considered mild (headache, gastrointestinal disorders, pharyngitis, fatigue, upper respiratory tract infection, cutaneous rash and reversible alterations in levels of serum transaminase) [33].

Although the main goal of root canal treatment is widely known to be to reduce the bacteria load and their by-products that contribute to the perpetuation of apical periodontitis [34, 35], promising results have been shown with the use of anti-inflammatory drugs and probiotics administered systemically or topically, and have demonstrated an advance in accelerating bone healing or preventing bone loss in apical periodontitis [8–10]. In this study, a histological and histometric evaluation was performed to understand the signaling pathways and mediators involved in bone resorption. This method is commonly used to determine periapical bone loss in induced apical periodontitis [10, 12] but this analysis could be improved through tomographic volumetric measurements [36–38]. This is one limitation of the present study, which should be better explored in future studies. Another limitation is regarding to the animal model, once the use of a mouse model where the root canal treatment was not performed may have impacted in the results observed, considering that the persistence of infection contributes to apical periodontitis development [1]. Moreover, in the search for

the ideal medication to control and reverse the apical periodontitis, the positive and negative aspects of the medicaments used should be evaluated, considering that it impacts on the choice of medicament to be used in routine clinical procedures.

Recently, Petean et al. (2021) [10] evaluated the efficacy of systemic use of selective and non-selective inhibitors of cyclooxygenase-2 enzymes in the treatment of experimental apical periodontitis induced by bacterial lipopolysaccharide (LPS) in vivo in a mouse model, and the results demonstrated promising results with the use of COX-2 inhibitor Celecoxib, which dampened osteoclastogenic signaling, and activity that suppressed bone resorption. Nevertheless, here it was demonstrated that inhibition of 5-LO by MK-886 is not suitable as a coadjuvant systemic treatment in clinical practice to prevent bone resorption in AP.

Conclusions

Systemic treatment with MK-886 exacerbated LPS-induced apical periodontitis in a murine model. Inhibition of 5-LO was, therefore, not a successful strategy to prevent bone resorption in apical periodontitis.

Acknowledgements

No applicable.

Author contributions

Conceptualization: IBFP, FWGGS; Formal analysis: IBFP, FWGGS; Methodology: IBFP, FWGGS; Data curation: IBFP, ACSS, RABS, MPL, FWGGS; Funding acquisition: IBFP, FWGGS; Writing-original draft: IBFP, RABS, MPL, FWGGS; Writing-review and editing: IBFP, ACSS, RABS, MPL, LABS, GPAC, MDSN, LHF, FWGGS; Supervision: FWGGS. All authors read and approved the final manuscript.

Funding

This work was supported by the São Paulo Research Foundation (FAPESP 2010/17611-4 to FWGGS, 2014/07125-6 to LHF and 2013/09595-7 to IBFP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq 303259/2020-5 to LHF).

Availability of data and materials

All the data generated during the study are available in the manuscript itself.

Declarations

Ethics approval and consent to participate

This manuscript was prepared in accordance with Brazilian Guideline for the Care and Use of Animals in Teaching or Scientific Research Activities regulated by the National Council for the Control of Animal experimentation (Law 11.794/2008). This study also conformed to ARRIVE (Animal Research: Reporting of In Vivo Experiments) guidelines and the animal study protocol was approved by the Ethical Committee at the University of São Paulo (São Paulo, Brazil, IRB approval number 12.1.60.53.8).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 9 August 2022 Accepted: 3 January 2023
Published online: 10 January 2023

References

- Nair PN. On the causes of persistent apical periodontitis: a review. *Int Endod J*. 2006;39(4):249–81.
- Petean IBF, Küchler EC, Soares IMV, et al. Genetic polymorphisms in RANK and RANKL are associated with persistent apical periodontitis. *J Endod*. 2019;45(5):526–31.
- Tibúrcio-Machado CS, Michelon C, Zanatta FB, Gomes MS, Marin JA, Bier CA. The global prevalence of apical periodontitis: a systematic review and meta-analysis. *Int Endod J*. 2021;54(5):712–35.
- Barbosa-Ribeiro M, Arruda-Vasconcelos R, Louzada LM, Dos Santos DG, Andreote FD, Gomes BPPA. Microbiological analysis of endodontically treated teeth with apical periodontitis before and after endodontic retreatment. *Clin Oral Investig*. 2021;25(4):2017–27.
- Chugal N, Mallya SM, Kahler B, Lin LM. Endodontic treatment outcomes. *Dent Clin North Am*. 2017;61(1):59–80.
- Bardini G, Casula L, Ambu E, Musu D, Mercadè M, Cotti E. A 12-month follow-up of primary and secondary root canal treatment in teeth obturated with a hydraulic sealer [published correction appears in *Clin Oral Investig*. 2021 Aug;25(8):5121]. *Clin Oral Investig*. 2021;25(5):2757–64.
- Siqueira JF Jr, Rôças IN. Clinical implications and microbiology of bacterial persistence after treatment procedures. *J Endod*. 2008;34(11):1291–1301. e3.
- Dessaune Neto N, Porpino MTM, Antunes HDS, et al. Pro-inflammatory and anti-inflammatory cytokine expression in post-treatment apical periodontitis. *J Appl Oral Sci*. 2018;26:e20170455.
- Cosme-Silva L, Dal-Fabbro R, Cintra LTA, et al. Systemic administration of probiotics reduces the severity of apical periodontitis. *Int Endod J*. 2019;52(12):1738–49.
- Petean IBF, Almeida-Junior LA, Arnez MFM, et al. Celecoxib treatment dampens LPS-induced periapical bone resorption in a mouse model. *Int Endod J*. 2021;54(8):1289–99.
- Ribeiro-Santos FR, Arnez MFM, de Carvalho MS, et al. Effect of non-steroidal anti-inflammatory drugs on pulpal and periapical inflammation induced by lipopolysaccharide. *Clin Oral Investig*. 2021;25(11):6201–9.
- Paula-Silva FW, Petean IB, da Silva LA, Faccioli LH. Dual role of 5-lipoxygenase in osteoclastogenesis in bacterial-induced apical periodontitis. *J Endod*. 2016;42(3):447–54.
- Ribeiro-Santos FR, Silva GGD, Petean IBF, et al. Periapical bone response to bacterial lipopolysaccharide is shifted upon cyclooxygenase blockage. *J Appl Oral Sci*. 2019;27:e20180641.
- Paula-Silva FW, Ribeiro-Santos FR, Petean IBF, et al. Root canal contamination or exposure to lipopolysaccharide differentially modulate prostaglandin E 2 and leukotriene B 4 signaling in apical periodontitis. *J Appl Oral Sci*. 2020;28:e20190699.
- Paula-Silva FW, Arnez MFM, Petean IBF, et al. Effects of 5-lipoxygenase gene disruption on inflammation, osteoclastogenesis and bone resorption in polymicrobial apical periodontitis. *Arch Oral Biol*. 2020;112:104670.
- Dixon RA, Diehl RE, Opas E, et al. Requirement of a 5-lipoxygenase-activating protein for leukotriene synthesis. *Nature*. 1990;343(6255):282–4.
- Okiji T, Morita I, Sunada I, Murota S. The role of leukotriene B4 in neutrophil infiltration in experimentally-induced inflammation of rat tooth pulp. *J Dent Res*. 1991;70(1):34–7.
- Torabinejad M, Cotti E, Jung T. Concentrations of leukotriene B4 in symptomatic and asymptomatic periapical lesions. *J Endod*. 1992;18(5):205–8.
- Cotti E, Torabinejad M. Detection of leukotriene C4 in human periradicular lesions. *Int Endod J*. 1994;27(2):82–6.
- Chen ZK, Lv HS, Jiang J. LTB4 can stimulate human osteoclast differentiation dependent of RANKL. *Artif Cells Blood Substit Immobil Biotechnol*. 2010;38(1):52–6.
- Dixit N, Wu DJ, Belgacem YH, Borodinsky LN, Gershwin ME, Adamopoulos IE. Leukotriene B4 activates intracellular calcium and augments human osteoclastogenesis. *Arthritis Res Ther*. 2014;16(6):496.
- Wu Y, Sun H, Yang B, Liu X, Wang J. 5-lipoxygenase knockout aggravated apical periodontitis in a murine model. *J Dent Res*. 2018;97(4):442–50.
- Graves DT, Oates T, Garlet GP. Review of osteoimmunology and the host response in endodontic and periodontal lesions. *J Oral Microbiol*. 2011. <https://doi.org/10.3402/jom.v3i0.5304>.
- Gillard J, Ford-Hutchinson AW, Chan C, et al. L-663,536 (MK-886) (3-[1-(4-chlorobenzyl)-3-t-butyl-thio-5-isopropylindol-2-yl]-2,2-dimethylpropanoic acid), a novel, orally active leukotriene biosynthesis inhibitor. *Can J Physiol Pharmacol*. 1989;67(5):456–64.
- Evans JF, Ferguson AD, Mosley RT, Hutchinson JH. What's all the FLAP about?: 5-lipoxygenase-activating protein inhibitors for inflammatory diseases. *Trends Pharmacol Sci*. 2008;29(2):72–8.
- Lopes DEM, Jabr CL, Dejana NN, et al. Inhibition of 5-lipoxygenase attenuates inflammation and BONE resorption in lipopolysaccharide-induced periodontal disease [published online ahead of print, 2017 Dec 5]. *J Periodontol*. 2017. <https://doi.org/10.1902/jop.2017.170210>.
- Le Bel M, Brunet A, Gosselin J. Leukotriene B4, an endogenous stimulator of the innate immune response against pathogens. *J Innate Immun*. 2014;6(2):159–68.
- Li YJ, Kukita A, Watanabe T, et al. Nordihydroguaiaretic acid inhibition of NFATc1 suppresses osteoclastogenesis and arthritis bone destruction in rats. *Lab Invest*. 2012;92(12):1777–87.
- Kang JH, Ting Z, Moon MR, et al. 5-Lipoxygenase inhibitors suppress RANKL-induced osteoclast formation via NFATc1 expression. *Bioorg Med Chem*. 2015;23(21):7069–78.
- Lorenzetti-Silva F, Arnez MFM, Thomé JPO, Carvalho MS, Carvalho FK, Queiroz AM, Faccioli LH, Paula-Silva FW. Leukotriene B4 Loaded in Microspheres Inhibits Osteoclast Differentiation and Activation. *Braz Dent J*. 2022;33(5):35–45.
- Coffey MJ, Phare SM, Peters-Golden M. Prolonged exposure to lipopolysaccharide inhibits macrophage 5-lipoxygenase metabolism via induction of nitric oxide synthesis. *J Immunol*. 2000;165(7):3592–8.
- Sforzini JM, Nunes GA, Missima F, Sá-Nunes A, Faccioli LH. Effect of a leukotriene inhibitor (MK886) on nitric oxide and hydrogen peroxide production by macrophages of acutely and chronically stressed mice. *J Pharm Pharmacol*. 2007;59(9):1249–54.
- Erdem SB, Nacaroglu HT, Unsal Karkiner CS, Gunay I, Can D. Side effects of leukotriene receptor antagonists in asthmatic children. *Iran J Pediatr*. 2015;25(5):e3313.
- Siqueira JF Jr. The defense: host-pathogen interactions and immune response. In: Siqueira JF, editor. *Treatment of endodontic infections*. 1st ed. London: Quintessence Publishing; 2011. p. 41–62.
- Siddique R, Nivedhitha MS. Effectiveness of rotary and reciprocating systems on microbial reduction: a systematic review. *J Conserv Dent*. 2019;22(2):114–22.
- Aksoy U, Savtekin G, Şehirli AÖ, Kermeoğlu F, Kalender A, Özkayalar H, Sayiner S, Orhan K. Effects of alpha-lipoic acid therapy on experimentally induced apical periodontitis: a biochemical, histopathological and micro-CT analysis. *Int Endod J*. 2019;52:1317–26.
- Chen S, Lei H, Luo Y, Jiang S, Zhang M, Lv H, Cai Z, Huang X. Micro-CT analysis of chronic apical periodontitis induced by several specific pathogens. *Int Endod J*. 2019;52:1028–39.
- Justo MP, Cardoso CBM, Cantiga-Silva C, de Oliveira PHC, Sivieri-Araújo G, Azuma MM, Ervolino E, Cintra LTA. Curcumin reduces inflammation in rat apical periodontitis. *Int Endod J*. 2022;55(1):1241–51.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.